



# Developmental neurotoxicity of industrial chemicals

P Grandjean, PJ Landrigan

Neurodevelopmental disorders such as autism, attention deficit disorder, mental retardation, and cerebral palsy, and autism are common, costly, and can cause lifelong disability. Their causes are mostly unknown. A few industrial chemicals (eg, lead, methylmercury, polychlorinated biphenyls [PCBs], arsenic, and toluene) are recognised causes of neurodevelopmental disorders and subclinical brain dysfunction. Exposure to these chemicals during early fetal development can cause brain injury at doses much lower than those affecting adult brain function. Recognition of these risks has led to evidence-based programmes of prevention, such as elimination of lead additives in petrol. Although these prevention campaigns are highly successful, most were initiated only after substantial delays. Another 200 chemicals are known to cause clinical neurotoxic effects in adults. Despite an absence of systematic testing, many additional chemicals have been shown to be neurotoxic in laboratory models. The toxic effects of such chemicals in the developing human brain are not known and they are not regulated to protect children. The two main impediments to prevention of neurodevelopmental deficits of chemical origin are the great gaps in testing chemicals for developmental neurotoxicity and the high level of proof required for regulation. New, precautionary approaches that recognise the unique vulnerability of the developing brain are needed for testing and control of chemicals.

One in every six children has a developmental disability and in most cases these disabilities affect the nervous system.<sup>1</sup> The most common neurodevelopmental disorders include learning disabilities, sensory deficits, developmental delays, and cerebral palsy.<sup>1</sup> Some experts have reported that the prevalence of certain neurodevelopmental disorders—autism and attention deficit and hyperactivity disorder, in particular—might be increasing, but there are few data to sustain that position.<sup>2</sup> Treatment of these disorders is difficult, and the disabilities they cause can be permanent;<sup>3</sup> they are therefore very costly to families and to society.<sup>4–6</sup>

Evidence has been accumulating over several decades that industrial chemicals can cause neurodevelopmental damage and that subclinical stages of these disorders might be common. The possibility of a link between chemicals and widespread neurobehavioural changes was first raised by research showing that lead was toxic to the developing brain across a wide range of exposures.<sup>7–10</sup> That report was in accord with reports indicating that other environmental pollutants were also toxic to early brain development.<sup>11</sup> An expert committee from the US National Research Council concluded that 3% of developmental disabilities are the direct result of environmental exposure to such poisons, and that another 25% arise through interactions between environmental factors and individual genetic susceptibility.<sup>3</sup> These estimates were based on scarce information about neurotoxicity and could therefore underestimate the true prevalence of chemically-induced abnormalities.

Neurobehavioural damage caused by industrial chemicals is, in theory, preventable. An essential prerequisite to prevention is recognition of a chemical's ability to harm the developing brain. Knowledge that a chemical is neurotoxic can prompt efforts to restrict its use and to control exposure. Previous evidence-based programmes of exposure prevention, such as those directed against children's exposure to lead, have been

highly successful, although they were initiated after substantial delay.

The aim of this review is to characterise the vulnerability of the developing nervous system to chemical toxicity; to collate publicly available data for human neurotoxicity of industrial chemicals; to examine the possible extent of a developmental neurotoxicity pandemic; to describe the known consequences of developmental neurotoxicity for individuals and society; to examine the implications for human health of the dearth of toxicological information; and to consider prospects for prevention of exposure.

## Vulnerability of the developing brain

The developing human brain is inherently much more susceptible to injury caused by toxic agents than is the brain of an adult.<sup>12</sup> This susceptibility stems from the fact that during the 9 months of prenatal life, the human brain must develop from a strip of cells along the dorsal ectoderm of the fetus into a complex organ consisting of billions of precisely located, highly interconnected, and specialised cells. Optimum brain development requires that neurons move along precise


Published Online  
November 8, 2006  
DOI:10.1016/S0140-6736(06)69665-7

Institute of Public Health, University of Southern Denmark, Odense, Denmark (Prof P Grandjean MD), Department of Environmental Health, Harvard School of Public Health, Boston, MA, USA (P Grandjean), Department of Community Medicine (Prof PJ Landrigan MD) and Department of Pediatrics, Mount Sinai School of Medicine, New York, NY, USA (PJ Landrigan)

Correspondence to:  
Dr Philippe Grandjean,  
Department of Environmental Health, Harvard School of Public Health, Landmark Center 3E-110, 401 Park Drive, Boston, MA 02215, USA.  
pgrand@hsph.harvard.edu

## Search strategy and selection criteria

We identified industrial chemicals that have caused neurotoxic effects in man from the hazardous substances data bank of the US National Library of Medicine, supplemented by fact sheets by the US Agency for Toxic Substances and Disease Registry, and the integrated risk information system of the US Environmental Protection Agency. We searched for the terms "neurotoxic", "neurological", and "neuro". For all neurotoxic substances identified, we then used synonyms, commercial names, and CAS (chemical abstracts service) numbers to search PubMed, TOXNET, and TOXLINE to identify published data for developmental neurotoxicity. The primary search terms were "prenatal exposure delayed effects"[MeSH] and "neurotoxicity syndromes"[MeSH]. Secondary searches used combinations of "maternal exposure" and "maternal fetal exchange" with "developmental disabilities/chemically induced" and "neurotoxins", all with the limiters "all child: 0–18 years", "most recent 10 Years", "English", and "human". We also used references cited in the chosen articles.



**Figure 1: The effects of a neurotoxic chemical on a population over time**  
For identification of chemicals toxic to neurodevelopment, the first evidence dealt with adverse effects of high doses on the adult nervous system, and was followed by case reports and epidemiological evidence on developmental toxicity at successively lower doses, to which childhood populations of increasing magnitude were exposed. Recognition of inorganic lead, methylmercury, and polychlorinated biphenyls as neurotoxic followed this curve towards the right, and arsenic and toluene were later seen to match this curve. Documentation of most neurotoxicants is directed toward adults only and therefore many compounds remain far to the left on the timescale.

pathways from their points of origin to their assigned locations, that they establish connections with other cells, both nearby and distant, and that they learn to communicate with other cells via such connections.<sup>12–14</sup> All these processes have to take place within a tightly controlled time frame, in which each developmental stage has to be reached on schedule and in the correct sequence. Because of the extraordinary complexity of human brain development, windows of unique susceptibility to toxic interference arise that have no counterpart in the mature brain, or in any other organ. If a developmental process in the brain is halted or inhibited, there is little potential for later repair, and the consequences can therefore be permanent.<sup>12,14</sup>

During fetal development, the placenta offers some protection against unwanted chemical exposures, but it is not an effective barrier against environmental pollutants.<sup>15</sup> For example, many metals easily cross the placenta, and the mercury concentration in umbilical cord blood can be substantially higher than in maternal blood.<sup>16</sup> The blood-brain barrier, which protects the adult brain from many toxic chemicals, is not completely formed until about 6 months after birth.<sup>17</sup>

The human brain continues to develop postnatally, and the period of heightened vulnerability therefore extends over many months, through infancy and into early childhood. Although most neurons have been formed by the time of birth, growth of glial cells and myelination of axons continues for several years.<sup>13,14</sup>

The susceptibility of infants and children to industrial chemicals is further enhanced by their increased exposures, augmented absorption rates, and diminished

ability to detoxify many exogenous compounds, relative to that of adults.<sup>18,19</sup> Persistent lipophilic substances, including specific pesticides and halogenated industrial compounds, such as PCBs, accumulate in maternal adipose tissue and are passed on to the infant via breast milk, resulting in infant exposure that exceeds the mother's own exposure by 100-fold on the basis of bodyweight.<sup>20</sup>

### Recognition of neurotoxicity

Developmental neurotoxicity in children exposed to industrial chemicals is often first identified through recognition of obvious functional abnormalities after high-dose exposure that clearly caused poisoning. Good quality research later documented the presence of less striking, but nonetheless serious adverse effects at low doses of exposure (figure 1). This sequence of discovery led to the recognition that environmental poisons exert a range of adverse effects—some are clinically evident, but others can be discerned only through special testing and are not evident on standard examination, hence the term subclinical toxicity. The underlying idea is that there is a dose-dependent continuum of toxic effects, in which clinically obvious effects have subclinical counterparts.<sup>21</sup> A pandemic of subclinical neurotoxicity is therefore likely to be silent—ie, not apparent from standard health statistics.

The notion of subclinical toxicity originates from Landrigan<sup>7</sup> and Needleman<sup>8</sup> and their colleagues' pioneering work who showed that children's exposure to lead could cause reductions in intelligence and changes in behaviour even in the absence of clinically visible symptoms of lead toxicity. The subclinical toxicity of lead in children has subsequently been confirmed in prospective epidemiological studies.<sup>22,23</sup>

Parallel findings have been reported on some other industrial chemicals, but their number is small. About 80 000 chemicals are registered for commercial use with the US Environmental Protection Agency, and 62 000 were already in use when the Toxic Substances Control Act was enacted in the USA in 1977.<sup>24</sup> The situation is similar in the EU, where 100 000 chemicals were registered in 1981.<sup>25</sup> The full extent to which these chemicals contribute to neurodevelopmental disorders and subclinical neurotoxicity is still unknown.

### Neurotoxic agents

#### Identification

Studies in animals support the notion that a wide range of industrial chemicals can cause developmental neurotoxicity at low doses that are not harmful to mature organisms.<sup>26,27</sup> Such injury seems to result in permanent changes in brain function that might become detectable only when the animal reaches maturity. Because developmental neurotoxicity might not be apparent from routine toxicology tests,<sup>28</sup> identification of neurotoxic chemicals often rests on clinical and epidemiological data.

To identify environmental chemicals that are toxic to the human brain, we searched the hazardous substances data bank of the US National Library of Medicine, where substances are listed with their adverse effects in man. We checked the completeness of this list against other data sources and with a previous review of published data for clinical toxicity.<sup>29</sup> The panel shows the industrial chemicals known to be neurotoxic in man. We have excluded drugs, food additives, microbial toxins, and snake venoms and similar biogenic substances. This list excludes chemicals that have proved neurotoxic solely in laboratory animals, for which no systematic list exists. We mainly include acute poisons that have caused serious accidents or have been used in suicide attempts, Neurotoxins that mainly cause chronic or delayed disease are likely to be underrepresented.<sup>29</sup> The largest groups of identified compounds are metals, solvents, and pesticides, but other chemicals with less documentation could have unrecognised effects. The list therefore should not be regarded as comprehensive.

Many substance names (see panel) were used for searches of published data for developmental neurotoxicity. On the basis of our critical review, the few known chemicals causing neurodevelopmental abnormalities are highlighted the panel. Many more chemicals that we have not listed are known to be neurodevelopment poisons in laboratory animals,<sup>27</sup> but no data about their potential toxic effects on human brain development are available.

### Lead

The neurotoxic effects of lead in adults were known in Roman times, but a report from Australia 100 years ago was the first description of epidemic lead poisoning in young children; the source of the outbreak was traced to ingestion of lead-based paint by children playing on verandas with peeling paint.<sup>30</sup> Further reports of childhood lead poisoning from the USA and Europe followed. Lead poisoning was at that time thought to be an acute illness, from which a child either recovered or died. Long-term sequelae were first documented in the 1940s, when 19 of 20 survivors of acute poisoning were noted to have severe learning and behavioural problems.<sup>31</sup>

Despite those early paediatric warnings, the largely unchecked use of lead in petrol, paints, ceramic glazes, and many other products through much of the twentieth century caused continued risk of lead poisoning. During the 1970s, widespread subclinical neurobehavioural deficits, including problems with concentration, memory, cognition, and behaviour, were documented in asymptomatic children with raised blood-lead concentrations.<sup>7,8</sup> Spurred by recommendations issued by the European Regional Office of WHO, studies were initiated in many countries; the results corroborated the previous conclusions.<sup>32</sup>

As a result of accumulating evidence, many sources of lead exposure became controlled, although not all sources,

#### Panel: Chemicals (n=201) known to be neurotoxic in man

##### Metals and inorganic compounds

- Aluminum compounds
- \*Arsenic and arsenic compounds
- Azide compounds
- Barium compounds
- Bismuth compounds
- Carbon monoxide
- Cyanide compounds
- Decaborane
- Diborane
- Ethylmercury
- Fluoride compounds
- Hydrogen sulphide
- \*Lead and lead compounds
- Lithium compounds
- Manganese and manganese compounds
- Mercury and mercuric compounds
- \*Methylmercury
- Nickel carbonyl
- Pentaborane
- Phosphine
- Phosphorus
- Selenium compounds
- Tellurium compounds
- Thallium compounds
- Tin compounds

##### Organic solvents

- Acetone
- Benzene
- Benzyl alcohol
- Carbon disulphide
- Chloroform
- Chloroprene
- Cumene
- Cyclohexane
- Cyclohexanol
- Cyclohexanone
- Dibromochloropropane
- Dichloroacetic acid
- 1,3-Dichloropropene
- Diethylene glycol
- N,N-Dimethylformamide
- 2-Ethoxyethyl acetate
- Ethyl acetate
- Ethylene dibromide
- Ethylene glycol
- n-Hexane
- Isobutyronitrile
- Isophorone
- Isopropyl alcohol

(Continues on next page)

(Continued from previous page)

- Isopropylacetone

- Methanol
- Methyl butyl ketone
- Methyl cellosolve
- Methyl ethyl ketone
- Methylcyclopentane
- Methylene chloride
- Nitrobenzene
- 2-Nitropropane
- 1-Pentanol
- Propyl bromide
- Pyridine
- Styrene
- Tetrachloroethane
- Tetrachloroethylene
- \*Toluene
- 1,1,1-Trichloroethane
- Trichloroethylene
- Vinyl chloride
- Xylene

#### Other organic substances

- Acetone cyanohydrin
- Acrylamide
- Acrylonitrile
- Allyl chloride
- Aniline
- 1,2-Benzenedicarbonitrile
- Benzonitrile
- Butylated triphenyl phosphate
- Caprolactam
- Cyclonite
- Dibutyl phthalate
- 3-(Dimethylamino)-propanenitrile
- Diethylene glycol diacrylate
- Dimethyl sulphate
- Dimethylhydrazine
- Dinitrobenzene
- Dinitrotoluene
- Ethylbis(2-chloroethyl)amine
- Ethylene
- Ethylene oxide
- Fluoroacetamide
- Fluoroacetic acid
- Hexachlorophene
- Hydrazine
- Hydroquinone
- Methyl chloride
- Methyl formate
- Methyl iodide
- Methyl methacrylate
- p-Nitroaniline
- Phenol

(Continues on next page)

(Continued from previous page)

- p-Phenylenediamine

and not in all countries. A 90% reduction in childhood blood-lead concentrations followed the termination of lead additives in petrol.<sup>33</sup> Now, research into lead neurotoxicity focuses on the shape of the dose-response curve at very low exposures that seem to cause surprisingly large functional decrements.<sup>22</sup> As convincing evidence was recognised, health agencies reduced the permissible concentration of lead in children's blood. However, up-to-date research<sup>22</sup> suggests that the current effects of lead exposure on human brain development could be even greater than previously thought.

#### Methylmercury

Toxic effects on the brain due to methylmercury were first established in men with occupational exposure.<sup>34</sup> The developmental toxicity of this organic mercury compound became evident in the 1960s in Minamata, Japan, where an epidemic of spasticity, blindness and profound mental retardation was seen in infants born to mothers who consumed fish from contaminated waters. After many years of clinical and experimental studies, the source proved to be mercury compounds released into Minamata Bay by a plastics plant.<sup>35</sup> Methylmercury accumulated and reached high concentrations in locally caught fish. Exposed adults, including mothers of poisoned children, were less seriously affected, if at all.<sup>36</sup> Similar outbreaks of profound neurodevelopmental disorders in the infants of seemingly unaffected mothers have arisen after maternal consumption during pregnancy of seed grain treated with methylmercury fungicides.<sup>37,38</sup> Studies of a serious poisoning incident in Iraq established a crude dose-response association between mercury concentrations in maternal hair and risk of neurological abnormalities in the children of the women.<sup>39</sup>

Recent studies have focused on prenatal exposures to reduced concentrations of methylmercury. They have examined populations with a high intake of seafood and freshwater fish with various degrees of methylmercury contamination. Prospective examination of a New Zealand cohort noted a three-point decrement in intelligence quotient (IQ) and changes in affect in children born to women with mercury concentrations in hair of greater than 6 µg/g.<sup>40</sup> A large prospective study in the Faroe Islands noted evidence of dose-related impairments in memory, attention, language, and visuospatial perception in exposed children.<sup>41</sup> A third prospective cohort study in the Seychelles provided some support for prenatal neurotoxicity after adjustment for postnatal exposures.<sup>42</sup> Several cross-sectional studies recorded significant associations between methylmercury exposure and neurobehavioral impairment in young children.<sup>43</sup>

The US National Academy of Sciences reviewed these studies and concluded that strong evidence exists for fetal neurotoxicity of methylmercury, even at low exposures.<sup>44</sup> These findings have led food safety

authorities to issue dietary advisories, and national and international agencies (with coordination from the UN Environment Programme) now seek to control and restrict mercury releases to the environment. Substantial reductions have already been achieved in mercury use and release from hospitals and incinerators.<sup>45</sup> A related substance, ethylmercury, has been widely used as a preservative in vaccines, but neurotoxic risk has not been documented.<sup>46</sup>

### Arsenic

Ingestion of arsenic-contaminated drinking water has long been recognised to cause peripheral neuropathy in adults.<sup>47</sup> Developmental neurotoxicity due to arsenic was reported in 1955 in Japan, where consumption of powdered milk contaminated with arsenic led to over 12 000 cases of poisoning and 131 deaths.<sup>48</sup> A follow-up study of three groups of adolescents born during the time of the milk contamination included one group that was fully breast-fed, one that was exposed to the tainted milk product, and one that received other supplements, but no tainted formula.<sup>49</sup> Compared with national rates, a ten-fold increase in mentally-retarded individuals were seen in the tainted milk group.<sup>48</sup> Poor school records, emotional disturbances, and abnormal or borderline electroencephalogram findings were also more common in the exposed group. Since these findings were initially reported in Japanese journals not easily available elsewhere,<sup>48,49</sup> they have often been overlooked, even in the most thorough risk assessments of environmental arsenic exposure.<sup>50,51</sup>

Arsenic is present in ground water worldwide, and industrial pollution is widespread. Cross-sectional studies of school-age children showed cognitive deficits associated with drinking water contamination<sup>52</sup> and raised urinary arsenic concentrations.<sup>53</sup> Similar results were obtained in children with arsenic exposure from a smelter.<sup>54</sup> Possible combined adverse effects on IQ caused by arsenic and manganese exposures was suggested by metal concentrations in hair in children living near a hazardous waste site.<sup>55</sup> Although evidence for subclinical neurodevelopmental neurotoxicity of arsenic is less well established than for lead and methylmercury, the data are consistent and fit with the high-exposure findings from Japan. Still, regulatory action does not emphasise the need to protect the developing brain against this neurotoxic substance.<sup>50,51</sup>

### Polychlorinated biphenyls

PCBs used to be widely used in electrical equipment as insulators. Human toxicity was first described from industrial exposures,<sup>56</sup> but neurological effects did not seem important. Developmental toxicity of PCBs was first seen in children exposed to high concentrations in two poisoning events in Asia, where cooking oil had been contaminated by PCBs and related substances during manufacturing. Prenatal exposure in one incident, in

- Phenylhydrazine
- Polybrominated biphenyls
- Polybrominated diphenyl ethers
- \*Polychlorinated biphenyls
- Propylene oxide
- TCDD
- Tributyl phosphate
- 2,2',2''-Trichlorotriethylamine
- Trimethyl phosphate
- Tri-o-tolyl phosphate
- Triphenyl phosphate

### Pesticides

- Aldicarb
- Aldrin
- Bensulide
- Bromophos
- Carbaryl
- Carbofuran
- Carbophenothion
- $\alpha$ -Chloralose
- Chlordane
- Chlordecone
- Chlorfenvinphos
- Chlormephos
- Chlorpyrifos
- Chlorthion
- Coumaphos
- Cyhalothrin
- Cypermethrin
- 2,4-D
- DDT
- Deltamethrin
- Demeton
- Dialifor
- Diazinon
- Dichlofenthion
- Dichlorvos
- Dieldrin
- Dimefox
- Dimethoate
- Dinitroresol
- Dinoseb
- Dioxathion
- Disulphoton
- Edifenphos
- Endosulphan
- Endothion
- Endrin
- EPN
- Ethiofencarb
- Ethion
- Ethoprop

(Continues on next page)

(Continued from previous page)



Taiwan, was associated with low birthweight, delayed developmental milestones, and lower IQs in comparison with unexposed siblings.<sup>57</sup> Exposed boys (but not girls)

- Fenitrothion
- Fensulphothion
- Fenthion
- Fenvalerate
- Fonofos
- Formothion
- Heptachlor
- Heptenophos
- Hexachlorobenzene
- Isobenzan
- Isolan
- Isoxathion
- Leptophos
- Lindane
- Merphos
- Metaldehyde
- Methamidophos
- Methidathion
- Methomyl
- Methyl bromide
- Methyl demeton
- Methyl parathion
- Mevinphos
- Mexacarbate
- Mipafox
- Mirex
- Monocrotophos
- Naled
- Nicotine
- Oxydemeton-methyl
- Parathion
- Pentachlorophenol
- Phorate
- Phosphamidon
- Phospholan
- Propaphos
- Propoxur
- Pyriminil
- Sarin
- Schradan
- Soman
- Sulprofos
- 2,4,5-T
- Tebupirimfos
- Tefluthrin
- Terbufos
- Thiram
- Toxaphene
- Trichlorfon
- Trichloronat

\*=substances that have been documented also to cause developmental neurotoxicity.

showed deficits in spatial reasoning. A follow-up study showed growth impairment, slow development, lack of endurance, clumsy movement, and very low IQs.<sup>58</sup> In a similar incident in Japan, neurological damage seemed less prominent than that of the Taiwan contamination.<sup>59</sup> Because of the mixed exposures, the specific contribution by PCB to these adverse effects cannot be determined.

Epidemiological studies of asymptomatic populations exposed prenatally to PCBs and related contaminants through maternal diet were done in the USA. Subclinical developmental deficits were shown in the most highly exposed of these children<sup>60,61</sup> and were associated, at age 11 years, with an average IQ score 6.2 points below that of children with lower exposures.<sup>62</sup> A Dutch cohort included 418 healthy infants and noted subclinical decrements on neonatal neurological examination and in subsequent developmental tests related to increased PCB exposures.<sup>63</sup> Continued follow-up of this cohort suggested that the effects could be modified or masked with age, but were still detectable at age 9 years.<sup>64</sup> Results from a German cohort were in accord with these findings and also suggested that postnatal PCB exposure from breastfeeding contributes to cognitive deficits.<sup>65</sup> A possible mechanism through which PCBs injure the developing brain is by interference with maternal thyroid function,<sup>13,66</sup> which might not harm adult brain functions. Although PCB manufacture has been banned in most nations, and exposures are decreasing, exposures at currently prevalent concentrations could still cause developmental neurotoxicity.<sup>67</sup>

### Solvents

Solvent neurotoxicity in adults is well known from acute poisoning cases and from occupational studies.<sup>68</sup> Ethanol is such a solvent. Intermittent, low-level exposures produce mild inebriating effects, but do not lead to irreversible damage. However, heavy, long-term ethanol intake in adults can lead to serious injury, including Wernicke's syndrome but because such exposures are voluntary, ethanol is not included in the panel.

Fetal alcohol syndrome is qualitatively different from the syndrome in adults. It was originally described in infants of mothers with a serious drinking habit, and involves cognitive and behavioural deficits and changes in facial features. Permanent neurotoxic damage in the mother is not a prerequisite for irreversible effects in the child.<sup>69</sup> At low consumption, subtle but permanent neurotoxicity, including decreased IQ scores, has been seen.<sup>70</sup> Effects of alcohol on the fetus could be enhanced by specific genetic polymorphisms.<sup>71</sup>

Less reliable documentation is available for other solvents widely used in industry. Because of its anaesthetic effects, toluene has been abused by sniffing, and case reports have reported that infants of mothers who sniffed toluene in pregnancy had abnormally low scores on developmental tests and showed delayed development of speech and motor function.<sup>72-74</sup> Additional evidence of cognitive deficits

in children comes from small studies of mothers who reported occupational exposure to solvents, including toluene, during pregnancy. [A: what were the effects on the children here?]<sup>75-77</sup> The women were apparently exposed within permissible workplace limits aimed at prevention of neurotoxicity in the workers themselves. However, these studies do not allow any definite conclusions on the specific hazard and the nature of dose-response associations for developmental neurotoxicity.

### Pesticides

More than 600 pesticides are registered, and include insecticides, fungicides, and rodenticides. In the USA alone, about 500 million kg [A: we use SI units is conversion ok?] are applied yearly. Acute pesticide neurotoxicity is well known from occupational exposure studies, poisoning events, and suicide data;<sup>29</sup> such neurotoxicity is often caused by cholinesterase inhibition by organophosphates.

Developmental neurotoxicity was suggested by an anthropological study of two similar groups of asymptomatic, Yaqui children aged 4–5 years in Mexico.<sup>78</sup> Those with high exposure to a mix of pesticides, including organophosphates, had diminished short-term memory, hand-eye coordination, and drawing ability, whereas unexposed children of the same tribe showed normal development.<sup>78</sup> Likewise, preschool children from agricultural communities in the USA showed poorer performance on motor speed and latency than did those of urban communities.<sup>79</sup> Ecuadorean schoolchildren, whose mothers had been exposed to organophosphates and other pesticides from working in greenhouses during pregnancy, showed visuospatial deficits compared with their unexposed peers.<sup>80</sup> Current pesticide exposure, measured by urinary excretion of organophosphate metabolites, was associated with delays in the children's simple reaction times.<sup>80</sup> Acute exposure of American children to the organophosphate pesticide, methyl parathion, was associated with persistent problems in short-term memory and attention span.<sup>81</sup> Prospective epidemiological studies of infants exposed prenatally to the organophosphate, chlorpyrifos, recorded significant decreases in head circumference and birthweight and slowing of reflexes.<sup>82-84</sup> Small head circumference, a risk factor for neurodevelopmental disorders, was seen only in exposed infants who were born to mothers with low expression of PON1, an esterase involved in organophosphate detoxification.<sup>83</sup> The effect of chlorpyrifos on bodyweight disappeared after introduction of a ban on residential use.<sup>84</sup> Although organophosphates can undoubtedly cause developmental neurotoxicity, the data are insufficient to determine the potential hazard to the developing brain posed by individual compounds among the dozens of organophosphates in use worldwide.

### Emerging neurotoxic substances

Documentation of developmental effects in human beings for the other compounds listed in the panel is

poor. However, three obvious candidate substances deserve particular attention, including two that have not seemed to cause neurotoxicity in adults.

### Manganese

Manganese neurotoxicity in adults has been well documented in occupationally exposed populations; parkinsonism is the classic clinical feature, and subclinical neurotoxicity has also been reported.<sup>85</sup> Concerns about the developmental neurotoxicity of manganese have emerged because the organic manganese compound methylcyclopentadienyl manganese tricarbonyl has been added to petrol as an antiknock agent in Australia and Canada and could be used in the USA and elsewhere in the future. Manganese can also be present in drinking water. In a prospective study of 247 births in Paris, France,<sup>86</sup> high manganese concentrations in cord blood were associated with impaired neurobehavioural development, especially on the Brunet-Lezine scales at age 9 months and the McCarthy scales at 3 years. At age 6, no association was seen but only 100 of the original children participated.<sup>86</sup> Community exposures to manganese released into the environment by combustion of methylcyclopentadienyl manganese tricarbonyl,<sup>87</sup> exposures from a toxic waste site in the USA,<sup>85</sup> and from contaminated drinking water in Bangladesh<sup>88</sup> have been associated with subclinical neurological impairment in children.

### Fluoride

Fluoride can cause neurotoxicity in laboratory animals,<sup>89</sup> but is not shown in the panel as a substance proven to be neurotoxic in man. It exists in drinking water as a natural contaminant, but the concentration is dependent on local geological circumstances. In rural communities in China, high fluoride concentrations in well water might cause skeletal abnormalities. In one such community, 222 children aged 8–13 years showed significantly worse IQs than 290 unexposed controls.<sup>90</sup> Parallel results were obtained in a smaller study of 118 children of similar age.<sup>91</sup> Another study of 477 schoolchildren from 22 villages suggested that both increased water fluoride concentrations and very low concentrations were associated with IQ deficits, compared with children exposed to normal concentrations (below 1 mg/L).<sup>92</sup> The reports did not thoroughly consider possible confounders, but do suggest that further in-depth studies be undertaken.

### Perchlorate

This chemical is not known as neurotoxic to adults. It is a widespread contaminant of ground water in the USA from the use of ammonium perchlorate as a solid-fuel propellant for rockets and missiles.<sup>93</sup> The thyroid is the primary site of perchlorate toxicity, and iodine uptake by the thyroid is blocked. Abnormal brain development, as a consequence of inhibition of maternal thyroid function,<sup>13,66</sup> is the major potential effect of perchlorate exposure.<sup>93</sup>

Because the available evidence is uninformative as to neurobehavioral toxicity, drinking water standards for perchlorate are set at levels to protect adults and do not include child-protective safety factors.

### Effects of developmental neurotoxicity

The five substances recognised as causes of developmental neurotoxicity show similar patterns in the development of scientific documentation of their risks. This pattern of discovery started in each instance with recognition of adult neurotoxicity, typically in people with occupational exposure and of episodes of acute, high-dose poisoning in children to the next stage was the accumulation of epidemiological evidence of neurobehavioural deficits in children with prenatal exposures at concentrations that are not toxic to adults (figure 1). For lead, methylmercury, and PCBs, widespread subclinical neurotoxicity has been documented internationally, yet the full implications of exposure to arsenic and toluene are unclear. For most substances listed in the panel, only neurotoxicity in adults has been documented.

The combined evidence suggests that neurodevelopmental disorders caused by industrial chemicals has created a silent pandemic in modern society. Although these chemicals might have caused impaired brain development in millions of children worldwide, the profound effects of such a pandemic are not apparent from available health statistics. Additionally, as shown by this Review only a few chemical causes have been recognised so the full effects of our industrial activities could be substantially greater than recognised at present.

As is shown by the evidence for inorganic lead, globally increased exposures have been responsible for erosion of cognitive skills with subclinical, but permanent decreases in IQ. Additionally, this neurotoxic chemical produces lifelong changes in behaviour with shortened attention span, increased impulsivity, heightened aggressiveness, slowed motor coordination, and impaired memory and language skills. The consequences are increased likelihood of school failure, diminished economic productivity, and possibly increased risk of antisocial and criminal behaviour.<sup>94</sup> The most striking of these effects occur at the extremes of exposure; in highly exposed children, almost none had above average function, whereas the number with obvious deficits increased greatly.<sup>95</sup> The most severely affected individuals will probably need special education and will also be less likely than their peers to pursue productive career options. A study of adults who were exposed to excess lead as children revealed that they were much less successful in life than those from a less exposed comparison group.<sup>96</sup>

The consequences of a pandemic of developmental neurotoxicity extend beyond descriptive data for incidence and prevalence of clinically diagnosed disorders.<sup>1,3</sup> Increased risk of Parkinson's disease<sup>97</sup> or other neurodegenerative diseases<sup>98</sup> is a further potential consequence

of the pandemic. Thus, early subclinical chemical injury has been postulated to silently kill a fraction of the cells needed to sustain brain function in later life (eg, in the substantia nigra). These latent impairments cause no symptoms in childhood, but could be unmasked during the natural neuronal attrition associated with ageing.<sup>99,100</sup>

The wide extent of human exposure to pollutants is now becoming apparent after systematic collection of data for the amount of these substances present in the environment and in human tissues.<sup>101</sup> Even then, recognition of causal associations could be difficult because exposures vary with time, more than one substance could have an effect, individual vulnerability varies, and there are other factors that can bias epidemiological studies toward the null hypothesis, especially when the outcome might be unrecognised for several years, or even decades.<sup>102</sup>

The population at risk of subclinical neurotoxicity from industrial chemicals is very large. Almost all children born in industrialised countries between 1960 and 1980 were exposed to substantial amounts of lead from petrol that could have reduced the number of children with far above average intelligence (IQ scores above 130 points) by over 50% and might likewise have increased the number with IQ scores below 70.<sup>95</sup> In the USA alone, the aggregate population of children at risk of exposure to airborne lead at that time was about 100 million. In this period, the resulting economic costs are estimated to have ranged from US\$110 billion to \$319 billion in each year's birth cohort.<sup>103</sup> Most of these costs were related to the diminished economic productivity that resulted over the exposed children's entire lifetimes from wide-scale reductions in intelligence. Today the costs of lead poisoning are estimated to be \$43 billion in each birth cohort in the USA,<sup>5</sup> whereas the costs of prenatal methylmercury toxicity are estimated to amount to \$8.7 billion yearly (range, \$2.2–43.8 billion).<sup>6</sup> Diminished economic productivity remains the main source of these costs. Because of the absence of dose-response associations for other neurotoxic compounds, the total costs are unknown.

The effect of chemical neurotoxicity extends beyond the industrially developed nations. Toxic chemicals, such as highly dangerous pesticides that are banned in industrialised countries, are exported to developing societies, where environmental and occupational standards are often weak or at least poorly enforced.<sup>104</sup> The consequences are largely unreported.

### Prevention

A pandemic of neurodevelopmental toxicity caused by industrial chemicals is, in theory, preventable. Testing of new chemicals before allowing them to be marketed is a highly efficient means to prevent toxicity, but has been required only in recent years. Of the thousands of chemicals used in commerce, fewer than half have been subjected to even token laboratory testing for toxicity




testing.<sup>24</sup> Nearly 3000 of these substances are produced in quantities of almost 500 000 kg every year, but for nearly half these high-volume chemicals no basic toxicity data are publicly available, and 80% have no information about developmental or paediatric toxicity.<sup>24</sup> Although new chemicals must be tested more thoroughly, access to these data can be restricted, because they could be claimed to constitute confidential business information. Absence of information about the neurotoxic potential of most industrial chemicals is therefore the main impediment to prevention of developmental disorders induced by neurotoxic pollutants. Accelerated testing of chemicals already in commerce is therefore essential. In the USA, a legal mandate to require testing was established in the Toxic Substances Control Act, but is largely unenforced.<sup>24</sup> In the EU, opportunity exists to require more extensive chemical testing through the REACH programme,<sup>25</sup> although the proposed legislation does not emphasise testing for developmental neurotoxicity as a primary objective.

Toxicity testing protocols for chemicals need to be expanded to include examination of neurobehavioural functions. Present test protocols rely mainly on crude indices, such as brain weight and gross morphology.<sup>105,106</sup> There is a risk that abbreviated protocols used for toxicity screening will overlook neurodevelopmental toxicity, and further testing could erroneously be thought unnecessary. Procedures for functional appraisal are available,<sup>105</sup> and a harmonised protocol for assessment of developmental neurotoxicity was developed under OECD auspices in 1999,<sup>106</sup> although a revision is still under review.

The number of chemicals that can cause neurotoxicity in laboratory studies probably exceeds 1000, which is far more than the estimated 200 that have caused documented human neurotoxicity. However, in the absence of systematic testing,<sup>28</sup> the true extent of the neurotoxic potential of industrial chemicals is unknown. The physiology of brain development<sup>12-14</sup> and experimental evidence<sup>14,26,27</sup> suggest that developmental neurotoxicity is likely for all of them, except perhaps for some of the compounds that require metabolic transformation to become neurotoxic, in which immature metabolism may provide some degree of protection.<sup>19,107</sup> The few substances proven to be toxic to human neurodevelopment should therefore be viewed as the tip of a very large iceberg (figure 2).

Large-scale, prospective epidemiological studies, such as birth cohorts from Europe<sup>108</sup> and the National Children's Study proposed in the USA, will be especially informative about early exposures and neurodevelopmental disorders.<sup>109</sup> Data from these investigations, especially when pooled internationally, will hopefully provide dose-response associations that can guide future disease prevention efforts. This research should move beyond repeated assessments of known neurotoxins to examine chemicals, whose toxicity is just beginning to be recognised. The substances listed in the panel, especially



**Figure 2: Diagram of the extent of knowledge of neurotoxic chemicals**  
Of the thousands of known chemicals, only a small fraction have been proven to cause developmental neurotoxicity in humans. Although this evidence does not represent the true potential for industrial chemicals to cause neurodevelopmental disorders, assessments of need for preventive measures nonetheless rely on that information.

those most prevalent in food, drinking water, and the environment, should provide a useful starting point. Nevertheless, these initiatives could take decades to generate the type of detailed documentation required for chemicals regulation.

The Food Quality Protection Act in the USA requires that pesticide standards be set at values that will protect infants against developmental toxicity. If testing data are not available, a child-protective safety factor should be used in standard settings. However, application of this factor has been uneven, and regulatory authorities need to recognise the vulnerability of prenatal brain development.

Prevention of neurodevelopmental disorders of chemical origin will need new approaches to control chemical exposures. The vulnerability of the human nervous system and its special susceptibility during early development suggest that protection of the developing brain should be a paramount goal of public health protection. The high level of proof needed for chemical control legislation has resulted in a slow pace of interventions to prevent exposures to lead and other recognised hazards. Instead, exposure limits for chemicals should be set at values that recognise the unique sensitivity of pregnant women and young children, and they should aim at protecting brain development. This precautionary approach, which is now beginning to be used in the EU, would mean that early indications of a potential for a serious toxic effect, such as developmental neurotoxicity, should lead to strict

regulation, which could later be relaxed, should subsequent documentation show less harm than anticipated.<sup>110</sup> As physicians, we should use prudence when counselling our patients, especially pregnant mothers, about avoidance of exposures to chemicals of unknown and untested neurotoxic potential.

#### Conflict of interest statement

P Grandjean has testified on behalf of the Natural Resources Defense Council in a court case in regard to mercury pollution from a chemical plant in Maine, USA. PJ Landrigan has testified on behalf of the State of Rhode Island, USA, in a lawsuit against the manufacturers of lead-based paint.

#### Acknowledgments

We thank Prof Katsuyuki Murata for providing and translating literature from Japanese sources, Anna Choi for the translations from Chinese, Jennifer Pintabone-Asetta for the database searches; and for financial support the Danish Medical Research Council, the US National Institute of Environmental Health Sciences (ES07384, ES09584, ES09797, ES11687), and the US Environmental Protection Agency (RD 83171101). These funding sources had no role in preparation of this review, and the contents of this paper are solely the responsibility of the authors and do not represent the official views of the US National Institute of Environmental Health Sciences or any other funding agency.

#### References

- Boyle CA, Decouffe P, Yeargin-Allsopp M. Prevalence and health impact of developmental disabilities in US children. *Pediatrics* 1994; **93**: 399–403.
- Rutter M. Incidence of autism spectrum disorders: changes over time and their meaning. *Acta Paediatr* 2005; **94**: 2–15.
- National Research Council. Scientific frontiers in developmental toxicology and risk assessment. Washington, DC: National Academy Press, 2000.
- Mendola P, Selevan SG, Gutter S, Rice D. Environmental factors associated with a spectrum of neurodevelopmental deficits. *Ment. Retard. Dev. D. R.* 2002; **8**: 188–197.
- Landrigan PJ, Schechter CB, Lipton JM, Fahs MC, Schwartz J. Environmental pollutants and disease in American children: Estimates of morbidity, mortality, and costs for lead poisoning, asthma, cancer, and developmental disabilities. *Environ Health Perspect* 2002; **110**: 721–08.
- Trasande L, Landrigan PJ, Schechter C. Public health and economic consequences of methyl mercury toxicity to the developing brain. *Environ Health Perspect* 2005; **113**: 590–96.
- Landrigan PJ, Whitworth RH, Baloh RW, Staehling NW, Barthel WF, Rosenblum BF. Neuropsychological dysfunction in children with chronic low-level lead absorption. *Lancet* 1975; **1**: 708–12.
- Needleman HL, Gunnoe C, Leviton A, Reed R, Peresie H, Maher C, Barrett P. Deficits in psychologic and classroom performance of children with elevated dentine lead levels. *N Engl J Med* 1979; **300**: 689–95.
- Baghurst PA, Robertson EF, McMichael AJ, Vimpani GV, Wigg NR, Roberts RR. The Port Pirie Cohort Study: lead effects on pregnancy outcome and early childhood development. *Neurotoxicology* 1987; **8**: 395–401.
- Dietrich KN, Krafft KM, Bornschein RL, Hammond PB, Berger O, Succop PA, Bier M. Low-level fetal lead exposure effect on neuro-behavioral development in early infancy. *Pediatrics* 1987; **80**: 721–30.
- Grandjean P, White RF. Developmental effects of environmental neurotoxicants. In: Tamburlini G, von Ehrenstein O, Bertollini R, eds. Children's health and environment. Environmental issue report No 29. Copenhagen: European Environment Agency, 2002, 66–78.
- Dobbing J. Vulnerable periods in developing brain. In: Davison AN, Dobbing J, eds. Applied Neurochemistry. Philadelphia: Davis, 1968, 287–316.
- Rodier PM. Developing brain as a target of toxicity. *Environ Health Perspect* 1995; **103** (Suppl 6): 73–6.
- Rice D, Barone S Jr. Critical periods of vulnerability for the developing nervous system: evidence from humans and animal models. *Environ Health Perspect* 2000; **108** (Suppl 3): 511–33.
- Andersen HR, Nielsen JB, Grandjean P. Toxicologic evidence of developmental neurotoxicity of environmental chemicals. *Toxicology* 2000; **144**: 121–127.
- Sakamoto M, Kubota M, Liu XJ, Murata K, Nakai K, Satoh H. Maternal and fetal mercury and n-3 polyunsaturated fatty acids as a risk and benefit of fish consumption to fetus. *Environ Sci Technol* 2004; **38**: 3860–63.
- Adinolfi M. The development of the human blood-CSF-brain barrier. *Dev Med Child Neurol* 1985; **27**: 532–37.
- National Research Council. Pesticides in the Diets of Infants and Children. Washington: National Academy Press, 1993.
- Ginsberg G, Hattis D, Sonawane B. Incorporating pharmacokinetic differences between children and adults in assessing children's risks to environmental toxicants. *Toxicol Appl Pharmacol* 2004; **198**: 164–83.
- Jensen AA, Slorach S, eds. Chemical contaminants in human milk. Boca Raton: CRC Press, 1991.
- Landrigan PJ. The toxicity of lead at low dose. *Br J Ind Med* 1989; **46**: 593–96.
- Lanphear BP, Hornung R, Khoury J, et al. Low-Level Environmental Lead Exposure and Children's Intellectual Function: An International Pooled Analysis. *Environ Health Perspect* 2005; **113**: 894–99.
- Bellinger DC. What is an adverse effect? A possible resolution of clinical and epidemiological perspectives on neurobehavioral toxicity. *Environ Res* 2004; **95**: 394–405.
- U.S. Environmental Protection Agency. Chemical Hazard Data Availability Study: What Do We Really Know About the Safety of High Production Volume Chemicals? Washington, DC: Office of Pollution Prevention and Toxics, 1998.
- Commission of the European Communities: White paper, Strategy for a Future Chemicals Policy (COM2001 88 final, 2001). Available at [http://europa.eu.int/comm/environment/chemicals/pdf/0188\\_en.pdf](http://europa.eu.int/comm/environment/chemicals/pdf/0188_en.pdf) (accessed, Aug 12, 2005);
- Tilson HA. Neurotoxicology risk assessment guidelines: developmental neurotoxicology. *Neurotoxicology* 2000; **21**: 189–94.
- Eriksson P. Developmental neurotoxicity of environmental agents in the neonate. *Neurotoxicology* 1997; **18**: 719–26.
- Claudio L, Kwa WC, Russell AL, Wallinga D. Testing methods for developmental neurotoxicity of environmental chemicals. *Toxicol Appl Pharmacol* 2000; **164**: 1–14.
- Kimbrough RD, Mahaffey KR, Grandjean P, Sandof[A: is this spelling correct?] SH, Ruttstein DD. Clinical effects of environmental chemicals: A software approach to etiologic diagnosis. New York: Hemisphere, 1989.
- Gibson, JL: A plea for painted railing and painted walls of rooms as the source of lead poisoning among Queensland children. *Aust Med Gazette* 1904; **23**: 149–53.
- Byers RK, Lord EE. Late effect of lead poisoning on mental development. *Am J Dis Child* 1943; **66**: 471–94.
- Winneke G, Brockhaus A, Ewers U, Kramer U, Neuf M. Results from the European multicenter study on lead neurotoxicity in children: implications for risk assessment. *Neurotoxicol Teratol* 1990; **12**: 553–59.
- Landrigan PJ: The worldwide problem of lead in petrol. *Bull World Health Organ* 2002; **80**: 768.
- Hunter D, Russell DS. Focal cerebral and cerebellar atrophy in a human subject due to organic mercury compounds. *J Neurol Neurosurg Psychiatr* 1954; **17**: 235–41.
- Harada M. Minamata disease: methylmercury poisoning in Japan caused by environmental pollution. *Crit Rev Toxicol* 1995; **25**: 1–24.
- Kitamura S, Miyata C, Tomita M, Date S. An epidemiological study on the unknown central nervous disorder in the Minamata district (in Japanese). *Kumamoto Med J* 1957; **32** (Suppl 1): 1–9.
- Elhassani SB. The many faces of methylmercury poisoning. *J Toxicol Clin Toxicol* 1982; **19**: 875–906.
- Pierce PE, Thompson JF, Likosky WH, Nickey LN, Barthel WF, Hinman AR. Alkyl mercury poisoning in humans. Report of an outbreak. *JAMA* 1972; **220**: 1439–42.
- Marsh DO, Myers GJ, Clarkson TW, et al. Dose-response relationship for human fetal exposure to methylmercury. *Clin Toxicol* 1981; **18**: 1311–18.
- Kjellström T, Kennedy P, Wallis S, et al. Physical and mental development of children with prenatal exposure to mercury from fish. Stage 2, interviews and psychological tests at age 6. (Report 3642) Stockholm, National Swedish Environmental Protection Board; 1989.

- 41 Grandjean P, Weihe P, White RF, et al. Cognitive deficit in 7-year-old children with prenatal exposure to methylmercury. *Neurotoxicol Teratol* 1997; **19**: 417–28.
- 42 Myers GJ, Davidson PW, Cox C, et al. Prenatal methylmercury exposure from ocean fish consumption in the Seychelles child development study. *Lancet* 2003; **361**: 1686–92.
- 43 Grandjean P, Cordier S, Kjellström T, Weihe P, Budtz-Jørgensen E. Health effects and risk assessments. In: Pirrone N, Mahaffey KR, eds. Dynamics of mercury pollution on regional and global scales: atmospheric processes and human exposures around the world. Norwell, MA: Springer, 2005, 499–523.
- 44 National Research Council. Toxicological effects of methylmercury. Washington: National Academy Press, 2000.
- 45 United Nations Environment Programme. Global Mercury Assessment. Geneva, Switzerland, 2002.
- 46 Hviid A, Stellfeld M, Wohlfahrt J, Melbye M. Association between thimerosal-containing vaccine and autism. *JAMA* 2003; **290**: 1763–66.
- 47 Landrigan PJ. Arsenic—state of the art. *Am J Ind Med* 1981; **2**: 5–14.
- 48 Yamashita N, Doi M, Nishio M, Hojo H, Tanaka M. Recent observations of Kyoto children poisoned by arsenic tainted 'Morinaga dry milk' [in Japanese]. *Nippon Eiseigaku Zasshi* 1972; **27**: 364–99.
- 49 Ohira M, Aoyama H. Epidemiological studies on the Morinaga Powdered Milk Poisoning Incident [in Japanese]. *Nippon Eiseigaku Zasshi* 1973; **27**: 500–31.
- 50 International Programme on Chemical Safety: Arsenic and arsenic compounds, second edition. Environmental Health Criteria 224. Geneva: World Health Organization, 2001.
- 51 National Research Council. Arsenic in drinking water. National Academy Press, 1999.
- 52 Tsai SY, Chou HY, The HW, Chen CM, Chen CJ. The effects of chronic arsenic exposure from drinking water on the neurobehavioral development in adolescence. *Neurotoxicology* 2003; **24**: 747–53.
- 53 Wasserman GA, Liu X, Parvez F, et al. Water arsenic exposure and children's intellectual function in Arahazar, Bangladesh. *Environ Health Perspect* 2004; **112**: 1329–33.
- 54 Calderon J, Navarro ME, Jimenez-Capdeville ME, et al. Exposure to arsenic and lead and neuropsychological development in Mexican children. *Environ Res* 2001; **85**: 69–76.
- 55 Wright RO, Amarasingwardena C, Woolf AD, Jim R, Bellinger DC. Neuropsychological correlates of hair arsenic, manganese, and cadmium levels in school-age children residing near a hazardous waste site. *Neurotoxicology* 2006; **27**: 210–16.
- 56 Smith AB, Schloemer J, Lowry LK, et al. Metabolic and health consequences of occupational exposure to polychlorinated biphenyls. *Br J Ind Med* 1982; **39**: 361–69.
- 57 Guo YL, Lambert GH, Hsu CC, Hsu MM. Yucheng: health effects of prenatal exposure to polychlorinated biphenyls and dibenzofurans. *Int Arch Occup Environ Health* 2004; **77**: 153–58.
- 58 Chen YCJ, Guo YL, Hsu CC, Rogan WJR. Cognitive development of Yu-Cheng ('Oil Disease') children prenatally exposed to heat-degraded PCBs. *JAMA* 1992; **268**: 3213–18.
- 59 Yoshimura T. Epidemiological study on Yusho babies to mothers who had consumed oil contaminated by PCB [in Japanese]. *Fukuoka Igaku Zasshi* 1974; **65**: 74–80.
- 60 Gladen BC, Rogan WJ, Hardy P, Thullen J, Tingelstad J, Tully M. Development after exposure to polychlorinated biphenyls and dichlorodiphenyl dichloroethene transplacentally and through human milk. *J Pediatr* 1988; **113**: 991–95.
- 61 Jacobson JL, Jacobson SW, Humphrey HEB. Effect of in utero exposure to polychlorinated biphenyls and related contaminants on cognitive functioning in young children. *J Pediatr* 1990; **116**: 38–45.
- 62 Jacobson JL, Jacobson SW. Intellectual impairment in children exposed to polychlorinated biphenyls in utero. *N Engl J Med* 1996; **335**: 783–89.
- 63 Patandin S, Lanting CI, Mulder PG, Boersma ER, Sauer PJ, Weisglas-Kuperus N. Effects of environmental exposure to polychlorinated biphenyls and dioxins on cognitive abilities in Dutch children at 42 months of age. *J Pediatr* 1999; **134**: 33–41.
- 64 Vreugdenhil HJ, Mulder PG, Emmen HH, Weisglas-Kuperus N. Effects of perinatal exposure to PCBs on neuropsychological functions in the Rotterdam cohort at 9 years of age. *Neuropsychology* 2004; **18**: 185–93.
- 65 Walkowiak J, Wiener JA, Fastabend A, et al. Environmental exposure to polychlorinated biphenyls and quality of the home environment: effects on psychodevelopment in early childhood. *Lancet* 2001; **358**: 1602–07.
- 66 Takser L, Mergler D, Baldwin M, de Grosbois S, Smargiassi A, Lafond J. Thyroid hormones in pregnancy in relation to environmental exposure to organochlorine compounds and mercury. *Environ Health Perspect* 2005; **113**: 1039–45.
- 67 European Food Safety Authority. Opinion of the CONTAM panel related to the presence of non dioxin-like polychlorinated biphenyls (PCB) in feed and food. [http://www.efsa.eu.int/science/contam/contam\\_opinions/1229\\_en.html](http://www.efsa.eu.int/science/contam/contam_opinions/1229_en.html) (accessed Aug 12, 2006).
- 68 White RF, Proctor SP. Solvents and neurotoxicity. *Lancet* 1997; **349**: 1239–43.
- 69 Sokol RJ, Delaney-Black V, Nordstrom B. Fetal alcohol spectrum disorder. *JAMA* 2003; **290**: 2996–9.
- 70 Streissguth AP, Barr HM, Sampson PD. Moderate prenatal alcohol exposure: Effects on child IQ and learning problems at age 7 and a half years. *Alcohol Clin Exp Res* 1990; **14**: 662–69.
- 71 Warren KR, Li TK. Genetic polymorphisms: impact on the risk of fetal alcohol spectrum disorders. *Birth Defects Res A Clin Mol Teratol* 2005; **73**: 195–203.
- 72 Hersh JH, Podruch PE, Rogers G, Weisskopf B. Toluene embryopathy. *J Pediatr* 1985; **106**: 922–27.
- 73 Arnold GL, Kirby RS, Langendoerfer S, Wilkins-Haug L. Toluene embryopathy: clinical delineation and developmental follow-up. *Pediatrics* 1994; **93**: 216–20.
- 74 Pearson MA, Hoyme HE, Seaver LH, Rimsza ME. Toluene embryopathy: delineation of the phenotype and comparison with fetal alcohol syndrome. *Pediatrics* 1994; **93**: 211–15.
- 75 Eskenazi B, Gaylord L, Bracken MB, Brown D. In utero exposure to organic solvents and human neurodevelopment. *Dev Med Child Neurol* 1988; **30**: 492–501.
- 76 Till C, Koren G, Rovet JF. Prenatal exposure to organic solvents and child neurobehavioral performance. *Neurotoxicol Teratol* 2001; **23**: 235–45.
- 77 Laslo-Baker D, Barrera M, Knittel-Keren D, et al. Child neurodevelopmental outcome and maternal occupational exposure to solvents. *Arch Pediatr Adolesc Med* 2004; **158**: 956–61.
- 78 Guillette EA, Meza MM, Aquilar MG, Soto AD, Garcia IE. An anthropological approach to the evaluation of preschool children exposed to pesticides in Mexico. *Environ Health Perspect* 1998; **106**: 347–53.
- 79 Rohlman DS, Arcury TA, Quandt SA, et al. Neurobehavioral performance in preschool children from agricultural and non-agricultural communities in Oregon and North Carolina. *Neurotoxicology* 2005; **26**: 589–98.
- 80 Grandjean P, Harari R, Barr DB, Debes F. Pesticide exposure and stunting as independent predictors of neurobehavioral deficits in Ecuadorian school children. *Pediatrics* 2006; **117**: 546–56.
- 81 Ruckart PZ, Kakolewski K, Bove FJ, Kaye WE. Long-term neurobehavioral health effects of methyl parathion exposure in children in Mississippi and Ohio. *Environ Health Perspect* 2004; **112**: 46–51.
- 82 Young JG, Eskenazi B, Gladstone EA, et al. Association between in utero organophosphate pesticide exposure and abnormal reflexes in neonates. *Neurotoxicology* 2005; **26**: 199–209.
- 83 Berkowitz GS, Wetmur JG, Birman-Deych E, et al. In utero pesticide exposure, maternal paraoxonase activity, and head circumference. *Environ Health Perspect* 2004; **112**: 388–91.
- 84 Whyatt RM, Rauh V, Barr DB, et al. Prenatal insecticide exposures and birth weight and length among an urban minority cohort. *Environ Health Perspect* 2004; **112**: 1125–32.
- 85 Pal PK, Samii A, Calne DB. Manganese neurotoxicity: a review of clinical features, imaging and pathology. *Neurotoxicology* 1999; **20**: 227–38.
- 86 Takser L, Mergler D, Hellier G, Sahuquillo J, Huel G. Manganese, monoamine metabolite levels at birth, and child psychomotor development. *Neurotoxicology* 2003; **24**: 667–74.
- 87 Mergler D, Baldwin M, Belanger S, et al. Manganese neurotoxicity, a continuum of dysfunction: results from a community based study. *Neurotoxicology* 1999 **20**: 327–42.

- 88 Wasserman GA, Liu X, Parvez F, et al. Water Manganese Exposure and Children's Intellectual Function in Araihasar, Bangladesh. *Environ Health Perspect* 2006; **114**: 124–29.
- 89 Mullenix PJ, Denbesten PK, Schunior A, Kernan WJ. Neurotoxicity of sodium fluoride in rats. *Neurotoxicol Teratol* 1995; **17**: 169–77.
- 90 Xiang Q, Liang Y, Chen L, et al. Effect of fluoride in drinking water on children's intelligence. *Fluoride* 2003; **36**: 84–94.
- 91 Lu Y, Sun ZR, Wu LN, Wang X, Lu W, Liu SS. Effect of high-fluoride water on intelligence in children. *Fluoride* 2000; **33**: 74–78.
- 92 Qin LS, Cui SY. The influence of drinking water fluoride on pupils' IQ as measured by Rui Wen's standard [in Chinese]. *Chinese J Control Epid Dis* 1990; **5**: 203–04.
- 93 National Research Council. Health Implications of Perchlorate Ingestion. Washington, DC: National Academy Press, 2005.
- 94 Needleman HL, McFarland C, Ness RB, Fienberg SE, Tobin MJ. Bone lead levels in adjudicated delinquents. A case control study. *Neurotoxicol Teratol* 2002; **24**: 711–17.
- 95 Needleman HL, Leviton A, Bellinger D. Lead-associated intellectual deficit (letter). *N Engl J Med* 1982; **306**: 367.
- 96 White RF, Diamond R, Proctor S, Morey C, Hu H. Residual cognitive deficits 50 years after lead poisoning during childhood. *Br J Ind Med* 1993; **50**: 613–22.
- 97 Langston W, Forno, LS, Tetrud J, Reeves AG, Kaplan JA, Karluk D. Evidence of active nerve cell degeneration in the substantia nigra of humans years after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine exposure. *J Ann Neurol* **46**: 598–605, 1999.
- 98 Calne DB, Eisen A, McGeer E, Spencer P. Alzheimer's disease, Parkinson's disease, and motoneuron disease: abiotrophic interaction between ageing and environment? *Lancet* 1986; **2**: 1067–70.
- 99 Grandjean P. Effects on reserve capacity, significance for exposure limits. *Sci Total Environ* 1991; **101**: 25–32.
- 100 Landrigan PH, Sonawane B, Butler RN, Trasande L, Callan R, Droller D: Early environmental origins of neurodegenerative disease in later life. *Environ Health Perspect* 2005; **113**: 1230–33.
- 101 Centers for Disease Control and Prevention. Third national report on Human Exposure to Environmental Chemicals. Atlanta: CDC, 2005. <http://www.cdc.gov/exposurereport> (accessed, Aug 12, 2005).
- 102 Axelson O. Negative and non-positive epidemiological studies. *Int J Occup Med Environ Health* 2004; **17**: 115–21.
- 103 Grosse SD, Matte TD, Schwartz J, Jackson RJ. Economic gains resulting from the reduction in children's exposure to lead in the United States. *Environ Health Perspect* 2002; **110**: 563–69.
- 104 Eddleston M, Karalliedde L, Buckley N, et al. Pesticide poisoning in the developing world—a minimum pesticides list. *Lancet* 2002; **360**: 1163–67.
- 105 Goldey ES, Tilson HA, Crofton KM. Implications of the use of neonatal birth weight, growth, viability, and survival data for predicting developmental neurotoxicity: a survey of the literature. *Neurotox Teratol* 1995; **17**: 313–32.
- 106 Hass U. Current status of developmental neurotoxicity: regulatory view. *Toxicol Lett* 2003; **140–141**: 155–59.
- 107 Scheuplein R, Charnley G, Dourson M. Differential sensitivity of children and adults to chemical toxicity. I. Biological basis. *Regul Toxicol Pharmacol* 2002; **35**: 429–47.
- 108 Kogevinas M, Andersen AM, Olsen J. Collaboration is needed to co-ordinate European birth cohort studies. *Int J Epidemiol* 2004; **33**: 1172–73.
- 109 Trasande L, Landrigan PJ. The National Children's Study: a critical national investment. *Environ Health Perspect* 2004 **112**: A789–90
- 110 Grandjean P. Implications of the precautionary principle for primary prevention and research. *Annu Rev Publ Health* 2004; **25**: 199–223.